


Vegetative Stages



True leaf — 4 cm



V-12



V-E




V-2



V-4

Stages of Sunflower Development

A.A. Schneiter, Professor
J.F. Miller, USDA-ARS
D.R. Berglund, Extension Agronomist



R-2



R-3

Reproductive Stages



R-1



R-2



R-3



R-3 Top View



R-4 Top View



R-5.1



R-5.5



R-5.9



R-6



R-7



R-8



R-9

NDSU

Extension Service

North Dakota State University
Fargo, North Dakota 58105

FEBRUARY 1998

Description of sunflower growth stages

The total time required for development of a sunflower plant and the time between the various stages of development depends on the genetic background of the plant and the growing environment. When determining the growth stage of a sunflower field, the average development of a large number of plants should be considered. This staging method can also be used for individual plants. The same system can be used for classifying either a single head or branched sunflower. In the case of branched sunflower, make determinations using only the main branch or head. In stages R7 through R9, use healthy, disease-free heads to determine plant development if possible, because some diseases can cause head discoloration.

	Stage	Description
Vegetative Emergence	VE	Seedling has emerged and the first leaf beyond the cotyledons is less than 4 cm long.
Vegetative Stages	V (number) (i.e.) V1 V2 V3 etc.,	These are determined by counting the number of true leaves at least 4 cm in length beginning as V1, V2, V3, V4, etc. If senescence of the lower leaves has occurred count leaf scars (excluding those where the cotyledons were attached) to determine proper stage.
Reproductive Stages	R1	The terminal bud forms a miniature floral head rather than a cluster of leaves. When viewed from directly above the immature bracts form a many-pointed star-like appearance.
	R2	The immature bud elongates 0.5 to 2.0 cm above the nearest leaf attached to the stem. Disregard leaves attached directly to the back of the bud.
	R3	The immature bud elongates more than 2.0 cm above the nearest leaf.
	R4	The inflorescence begins to open. When viewed from directly above immature ray flowers are visible.
	R5 (decimal) (i.e.) R5.1 R5.2 R5.3 etc.	This stage is the beginning of flowering. The stage can be divided into substages dependent upon the percent of the head area (disk flowers) that has completed or is in flowering. Ex. R5.3 (30%), R5.8 (80%) etc.
	R6	Flowering is complete and the ray flowers are wilting.
	R7	The back of the head has started to turn a pale yellow color.
	R8	the back of the head is yellow but the bracts remain green.
	R9	The bracts become yellow and brown. This stage is regarded as physiological maturity.

From Schneider, A.A., and J.F. Miller. 1981. Description of Sunflower Growth Stages. *Crop Sci.* 21:901-903.

For more information on this and other topics, see: www.ag.ndsu.edu



NDSU Extension Service, North Dakota State University of Agriculture and Applied Science, and U.S. Department of Agriculture cooperating. Duane Hauck, director, Fargo, N.D. Distributed in furtherance of the acts of Congress of May 8 and June 30, 1914. We offer our programs and facilities to all people regardless of race, color, national origin, religion, sex, disability, age, Vietnam era veterans status or sexual orientation; and are an equal opportunity employer.

2M-2-98, 1.5M-4-05

This publication will be made available in alternative format upon request to people with disabilities (701) 231-7881.